

CURRICULUM VITAE

Name **OCHIAI, Masayuki “Masa” MD, PhD**

Present Position

Associate Professor, Research Center for Environmental and Developmental Medical Sciences, Kyushu University, Fukuoka, JAPAN

Business Address

3-1-1, Maidashi, Higashi-ku, Fukuoka, 812-8582, JAPAN

FAX 092-642-5435

<https://pediatr.kyushu-u.ac.jp>

Academic Degree

1999 Doctor of Medicine (MD)

Graduate School of Medical Sciences, Kyushu University

2009 Doctor of Philosophy (PhD)

Department of Pediatrics, Graduate School of Medical Sciences, Kyushu University

Educations and Professional Background

1999-2000 Resident, Kyushu University Hospital

2000-2003 Resident,

Oita Prefectural Hospital / Fukuoka Children's Hospital / Kyushu Medical Center

2004-2007 Postgraduate School of Medicine, Kyushu University

2008-2017 Assistant Professor, Kyushu University Hospital

2018-2021 Associate Professor, Department of Perinatal and Pediatric Medicine,

Graduate School of Medical Sciences, Kyushu University

2022- Associate Professor, Research Center for Environmental and Developmental Medical Sciences, Kyushu University

Membership

Japan Pediatric Society

Japan Society of Perinatal and Neonatal Medicine (Advisor, Councilor)

Japan Society for Neonatal Health and Development (Director, Councilor)

The Japan Society of Obstetrical, Gynecological and Neonatal Hematology (Advisor, Councilor)

The Japanese Society of Child Neurology, The Japan Society of Human Genetics

The Japanese Society on Thrombosis and Hemostasis

Bibliography (Neonatology; Fundamental)

- 1 Sakemi Y, Nakashima T, Watanabe K, Ochiai M, Sawano T, Inoue H, Kawakami K, Isomura S, Yamashita H, Ohga S: Changing risk factors for postpartum depression in mothers admitted to a perinatal center. *Pediatr Neonatol.* 2023 May;64(3):319-326.
- 2 Matsushita Y, Sakai Y, Torio M, Inoue H, Ochiai M, Yasuoka K, Kurata H, Fujiyoshi J, Ichiyama M, Taguchi T, Kato K, Ohga S; Neonatal Research Network of Japan (NRNJ): Association of perinatal factors of epilepsy in very low birth weight infants, using a nationwide database in Japan. *J Perinatol.* 2019 Nov;39(11):1472-1479.
- 3 Kurata H, Ochiai M, Inoue H, Ichiyama M, Yasuoka K, Fujiyoshi J, Matsushita Y, Honjo S, Sakai Y, Ohga S; Neonatal Research Network of Japan: A nationwide survey on tracheostomy for very-low-birth-weight infants in Japan. *Pediatr Pulmonol.* 2019 Jan;54(1):53-60.
- 4 Kurata H, Ochiai M, Inoue H, Kusuda T, Fujiyoshi J, Ichiyama M, Wakata Y, Takada H: Inflammation in the Neonatal Period and Intrauterine Growth Restriction Aggravate Bronchopulmonary Dysplasia. *Pediatr Neonatol.* 2019 Oct;60(5):496-503.
- 5 Inoue H, Ochiai M, Sakai Y, Yasuoka K, Tanaka K, Ichiyama M, Kurata H, Fujiyoshi J, Matsushita Y, Honjo S, Nonaka K, Taguchi T, Kato K, Ohga S; Neonatal Research Network of Japan: Neurodevelopmental Outcomes in Infants With Birth Weight ≤ 500 g at 3 Years of Age. *Pediatrics.* 2018 Nov 16. pii: e20174286.
- 6 Inoue H, Ochiai M, Yasuoka K, Tanaka K, Kurata H, Fujiyoshi J, Matsushita Y, Suga S, Nonaka K, Taguchi T, Kato K, Ohga S; Neonatal Research Network of Japan (NRNJ): Early Mortality and Morbidity in Infants with Birth Weight of 500 Grams or Less in Japan. *J Pediatr.* 2017 Nov;190:112-117.e3.
- 7 Ochiai M, Matsushita Y, Inoue H, Kusuda T, Kang D, Ichihara K, Nakashima N, Ihara K, Ohga S, Hara T; Kyushu University High-Risk Neonatal Clinical Research Network, Japan: Blood Reference Intervals for Preterm Low-Birth-Weight Infants: A Multicenter Cohort Study in Japan. *PLoS One.* 2016 Aug 23;11(8):e0161439.
- 8 Ochiai M, Kurata H, Inoue H, Tanaka K, Matsushita Y, Fujiyoshi J, Wakata Y, Kato K, Taguchi T, Takada H: An Elevation of Serum Ferritin Level Might Increase Clinical Risk for the Persistence of Patent Ductus Arteriosus, Sepsis and Bronchopulmonary Dysplasia in Erythropoietin-Treated Very-Low-Birth-Weight Infants. *Neonatology.* 2016 Aug 23;111(1):68-75.
- 9 Ochiai M, Ichiyama M, Iwayama M, Sakai Y, Yoshida K, Hara T: Longitudinal study of very low birth weight infants until 9 years of age; attention deficit hyperactivity and autistic features are correlated with their cognitive functions. *Early Hum Dev.* 2015 Dec;91(12):783-6.
- 10 Ochiai M, Kinjo T, Takahata Y, Iwayama M, Abe T, Ihara K, Ohga S, Fukushima K, Kato K, Taguchi T, Hara T: Survival and neurodevelopmental outcome of preterm infants born at 22-24 weeks of gestational age. *Neonatology.* 2014; 105: 79-84.
- 11 Inoue H, Ohga S, Kusuda T, Kitajima J, Kinjo T, Ochiai M, Takahata Y, Honjo S, Hara T: Serum neutrophil gelatinase-associated lipocalin as a predictor of the development of bronchopulmonary dysplasia in preterm infants. *Early Hum Dev.* 2013; 89 (6): 425-9.
- 12 Ochiai M, Hikino S, Yabuuchi H, Nakayama H, Sato K, Ohga S, Hara T: A new scoring system for computed tomography of the chest for assessing the clinical status of bronchopulmonary dysplasia. *J Pediatr.* 2008; 130 (7): 2118-9.